

[0003] Models used to predict stock market activity have tended to be based on the assumption that price change activity follows a normal distribution.

[0005] These variations of market behavior from expected norms lead to the failure of the known models for market assessment and prediction.

25 Summary of the Invention

[0008] The invention is based on the proposition that if there are no undue outside influences on a financial market, and there is no collusion within the market, prices can be expected to be haphazardly pushed
35 about so as to oscillate around a mean in a manner

predicted by Brownian motion. A particle subject to Brownian motion is pushed around by the random motions of neighboring particles, and takes time Δt to move throughout a circular two-dimensional area of radius r ,
5 and can be expected to take $4\Delta t$ to cover the circular area of radius $2r$ (including the original area of radius r) because that area is four times the original area (an r -squared relationship). The same is true of, e.g., stock prices -- if prices move within a range Δp
10 during a time interval Δt , then during the ensuing $3\Delta t$ (for a total elapsed time of $4\Delta t$), prices should move erratically or haphazardly within a range of magnitude $2\Delta p$ (i.e., it takes four times as long to move twice Δp as it takes to move once Δp). If in fact the range of
15 price movement during the longer interval is smaller than twice what it was during the shorter interval, then there is some other influence, such as a congealed group of buyers or sellers, who may be acting independently or colluding, whose collective buying
20 and/or selling acts to push or constrain price in a particular direction, and the price movement is in a congestion situation. On the other hand, if the range of price movement during the longer interval is larger than twice what it was during the shorter interval,
25 then there is some other influence such as the aforementioned congealed group of buyers or sellers, and the price movement is in a trend situation. In a trend situation, the absolute magnitude of the price will reveal whether the trend is an upward trend or a
30 downward trend. Note that it is not necessary to use a longer period that is four times the shorter period; rather, any longer period can be used, and the range of expected movement is proportional to the square root of the ratio of the durations of the longer and shorter
35 periods. For example, if the longer period is nine times the shorter period, the expected range of

09736070 121300

movement in the longer period would be three times the range of movement in the shorter period.

[0009] This relationship should hold regardless of the absolute magnitude of the time periods involved.

5 Thus, using the four-to-one ratio, the movement over the last four minutes should be twice the movement that occurred in the one-minute interval between four minutes ago and three minutes ago, just as the movement over the last four hours should be twice the movement
10 that occurred in the one-hour interval between four hours ago and three hours ago and the movement over the last four days should be twice the movement that occurred in the one-day interval between four days ago and three days ago. By continually monitoring price
15 changes, one can continually compute, based on any time interval or intervals one chooses, whether the market is in a trend situation or a congestion situation. One can choose to monitor only the previous four minutes, or only the previous four hours, or both the previous
20 four minutes and the previous four hours, or days, or months, or years, or any other combination.

[0010] Instead of, or in addition to, monitoring the range of prices during an interval and treating that interval as the longer interval, going back and taking
25 the first quarter of that interval as the shorter interval, one could determine the range during an interval and treat that interval as the shorter interval. From that range measured during the shorter interval one could predict the expected range during
30 the ensuing longer interval of which the shorter interval is a part. For the remainder of the longer interval one could simply monitor the actual price at each moment during the longer interval, looking forward instead of waiting for the interval to be completed and
35 looking backward to calculate the actual range. If the price moves outside the expected range, then clearly

09736070 121300

the magnitude of the actual range is going to be larger than that of the expected range, and one can assume that a trend has begun without having to determine what the actual range is. If the actual price is within the
5 expected range, however, it may remain within a smaller range (indicating congestion), so in that case it would be necessary to wait until the longer interval is complete and determine the actual range.

[0011] As an alternative to the "square-root-of-
10 time" relationship just described, one could compare
price movements against a generated Brownian motion
standard. Such standards are increasingly common in
personal computers, in which they may be provided in
the form of a separate chip the output of which is
15 based, e.g., on fluctuations of chip temperature, and
are provided to facilitate encryption algorithms. If
such a standard is used, then the time intervals for
which price changes are measured can be arbitrary.

[0012] Because price ranging is subject to an
20 r-squared relationship (as discussed above), price
ranging can be compared to a planetary orbit, which is
also governed by an r-squared relationship (i.e.,
gravity), with the core force exerted by the star
around which the planet orbits. When a planet is
25 orbiting about a star with no influence other than the
inverse-square force of the star, the planetary orbit
will be circular. However, for known planets there is
always another influence, which is the net
gravitational force exerted by the rest of the
30 universe. Where there is such a second influence, the
planetary orbit becomes distorted into a familiar
elliptical shape, with the planet moving fastest when
it is near its star, slowest when it is furthest from
its star, and at its prior medium or undistorted speed
35 somewhere in between.

[0013] Following this analogy, price ranging can be considered as following an elliptical orbit, with the core influence stemming from the Brownian motion force exerted by the bulk of buying and selling -- i.e., the "star" around which price ranging (the "planet") orbits. The disturbing force is the congealed group of buyers or sellers trying to move the price concertedly in one direction (either up or down). This orbital analogy teaches us first that every trend must be followed by a congestion period -- and vice-versa -- because the orbit is closed, unless the outside influence is so great that the orbital ellipse opens up to a parabola or hyperbola, in which prices fly off one end, "crash," and do not automatically return (e.g., in October 1929). Moreover, by fitting price change data to an ellipse, one can determine the shape of the ellipse and where on the ellipse price ranging currently is. Using Kepler's Laws of planetary motion, which postulate that an orbiting planet will sweep out equal areas in equal times, and knowing how long the market price ranging took to move between the plotted points, one can determine when the market price ranging will move from its current condition to a different condition -- i.e., if market price ranging is in a trend, when it will move into a congestion period, and if market price ranging is in a congestion period when it will break out into a trend.

[0014] Although the invention has been described thus far with regard to market price data, it can be applied to other financial data, or to non-financial data such as biological or meteorological data.

[0015] Because time continues to move forward, while one is performing the calculations for a selected interval, the four-time-unit window is moving. Depending on the granularity chosen, one must soon begin the next calculation. For example, if one is

09736070-121300

calculating based on a four-hour window, one may want to sample that four-hour window every ten minutes, or every minute, or perhaps only once an hour. If the sampling is too infrequent, one may miss the turning
5 point between a trend and congestion, or vice-versa. But if the sample is too frequent, it may simply add to the computational burden with little gain of useful data. Thus, sampling the previous four hours every second may be too frequent, but sampling the previous
10 four hours only every two hours may not be frequent enough. Sampling every minute would appear to be a practical approach.

[0016] In any event, all of this continual recalculation of the four-time-unit data requires
15 significant computing resources. In accordance with another aspect of the invention, those computing resources are provided by all of those using the method of the invention. As part of a subscription to the data feed produced by the method, a subscriber
20 preferably would agree to accept a distributed portion of the data to be analyzed and to have his or her computer perform the analysis. One way that this could be done would be to have the analysis performed only while the subscriber's computer is idle, such as by a
25 program that operates when the computer is in "screen saver" mode. In one embodiment of such distributed computing system, a subscriber would have the option of whether or not to make his or her computer available, with those subscribers who make their computers
30 available receiving a discounted subscription rate as compared to those subscribers who do not make their computers available.

Brief Description of the Drawings

[0017] The above and other objects and advantages of the invention will be apparent upon consideration of the following detailed description, taken in
5 conjunction with the accompanying drawings, in which like reference characters refer to like parts throughout, and in which:

[0018] FIG. 1 is an example of Brownian motion of a particle in a shallow dish of liquid;

10 [0019] FIG. 2 is a graph comparing the ranging of stock prices to Brownian motion;

[0020] FIG. 3 is a graph showing greater ranging of stock prices than would be predicted by Brownian motion;

15 [0021] FIG. 4 is a graph showing less ranging of stock prices than would be predicted by Brownian motion;

[0022] FIG. 5 is a plot comparing the ranging of stock prices to a planetary orbit;

20 [0023] FIG. 6 is a schematic block diagram of an exemplary hardware system with which the invention can be used;

[0024] FIG. 7 is a graph comparing the ranging of stock prices to a Gaussian bell curve produced by a
25 Brownian motion standard;

[0025] FIG. 8 is a plot showing ranging of prices along the orbit plotted in FIG. 5; and

[0026] FIG. 9 is a representation of a preferred output display produced by the invention.

30 Detailed Description of the Invention

[0027] As described above, the present invention allows one to monitor changing data over time intervals, and to determine, from the change of the data during the interval, the present state of the data

09736070.121300

relative to normative value, and how the data are likely to change in the future or, more specifically, whether or not current patterns in the data can be expected to continue. The types of data for which the invention will operate are those types where a datum is pushed about in various directions by forces governed by a relationship based on the inverse power of the number of degrees of freedom, much like the motion of a small particle suspended in a thin layer liquid, being pushed about by invisible collisions with molecules of the liquid which, as a two-dimensional system, is governed by an inverse-square law (a three-dimensional system would be governed by an inverse-cube law). Thus, a stock price, whose measurements over time represent a two-dimensional system, does not change purely randomly, as has been believed, but moves based on "collisions" with orders from buyers and sellers who are buying and selling predominantly haphazardly.

[0028] The invention will now be explained with
20 reference to FIGS. 1-9.

[0029] FIG. 1 shows the movement of a particle 10 in a Petri dish 11 filled with a thin layer of liquid 12. In a first time interval Δt , the particle moves throughout the area of the circle 13 of radius r . The area of the circle 14 of radius $2r$ is four times the area of circle 13. Therefore, the time necessary for the particle 10 to move throughout the area of circle 14 is four times Δt , or $4\Delta t$. This longer time interval includes the initial time interval Δt (insofar as circle 13 is wholly within circle 14), and thus concludes $3\Delta t$ after the end of the initial time interval Δt .

[0030] FIG. 2 is a graph of stock price changes over time. Instead of circles of radii r and $2r$, what is
35 important is the range of price changes, between a minimum price during an interval and the maximum price

[illegible]

during that interval. For a first interval of duration Δt beginning at time t_0 and a second interval of duration $4\Delta t$ beginning at time t_0 , one would expect the range 20 between minimum and maximum prices during the longer interval to be twice the range 21 between minimum and maximum prices during the shorter interval, if the changes produced by buying and selling occurred in a purely Brownian motion fashion.

[0031] FIG. 3 is similar to FIG. 2, but range 30 is more than twice range 21. This is a situation where an outside force is tugging against the movement of price and distorting it from expanding at a uniform rate as depicted by the Brownian motion circle, and represents a price trend. In this particular case, the trend is upward, which can be seen from the magnitudes of the changes.

[0032] FIG. 4 is similar to FIGS. 2 and 3, but range 40 is less than twice range 21. This is a situation where an outside force is constraining the movement of price and distorting it from expanding at a uniform rate as depicted by the Brownian motion circle, and represents a price congestion condition.

[0033] Note that while the time intervals in FIGS. 2-4 are shown as being related by a multiple of 4, with an expected price range multiple of 2, the multiple of the time intervals can have any value, with the expected price range multiple being the square root of the multiple of the time intervals. Moreover, if the price changes are being compared to a generated Brownian motion standard, then any time interval can be used.

[0034] The graphs of FIGS. 2-4 can be represented as orbital models as discussed above. The condition of FIG. 2, in which the range of prices during the longer time interval is exactly twice the range during the shorter interval normally can be expected to exist for

003736070 124300

only very short times, as conditions move between the other states shown in FIGS. 3 and 4. However, if the condition of FIG. 2 were to persist, it would signify that price changes were purely Brownian, with no
5 congealed group of buyers or sellers acting to distort the inverse-square law effect. The abstract orbital representation of such a price-expansion condition would be a circle of arbitrary radius (not shown), representing an orbit around a single source of
10 inverse-square law force, with no disturbing force. [0035] The more common price-expansion situations, illustrated in FIGS. 3 and 4, can be analogized to non-circular elliptical planetary orbits as discussed above, with the core inverse-square law Brownian motion
15 "force" located at one focus of the ellipse in the role of the star, and the net effect of the distorting "force" resulting from the congealed group of buyers or sellers located at the second focus of the ellipse. The values calculated according to the present
20 invention represent velocity along the elliptical orbit. Such a non-circular elliptical orbit 50 is shown in FIG. 5, with the Brownian "star" 51 at one focus, and the net effect of the distorting forces acting at a second focus 55. Just as in planetary
25 mechanics, the orbiting body will sweep equal areas between itself and the "star" in equal times. Therefore, when the orbiting body is close to "star" 51, the orbital velocity will be high, and when it is far from "star" 51 the orbital velocity will be
30 low. The high-velocity portion 52 of orbit 50 represents conditions where the range of prices during the longer interval exceeds the range during the shorter interval by more than would be expected based on a "square root of time" relationship, while the low-
35 velocity portion 53 of orbit 50 represents conditions where the range of prices during the longer interval

09736070 121300

exceeds the range during the shorter interval by less than would be expected based on a "square root of time" relationship. The line 54 represents the points where the "square root of time" relationship is met exactly
5 as one moves from region 52 to region 53 or vice-versa. To a gross approximation, if only one point were to be measured -- i.e., only one comparison between a shorter interval and a longer interval was made, region 52 could be considered a price trend region, while
10 region 53 could be considered a price congestion region. However, few useful predictions could be made based on one point.

[0036] Somewhat better predictions could be made based on two points. If two consecutive points are
15 taken (i.e., two comparisons between respective long and short intervals), one can place current conditions in a particular quadrant of orbit 50. Thus, if two points are measured, both exceed the "square root of time" expectation, and the second is higher than the
20 first, then the measured phenomenon -- whether a stock market or some other phenomenon including a natural phenomenon -- is in quadrant I (assuming a clockwise orbit), meaning that it is in a trend condition and the trend is accelerating. If two points are measured,
25 both exceed the "square root of time" expectation, and the second is lower than the first, then the measured phenomenon is in quadrant II (assuming a clockwise orbit), meaning that it is in a trend condition but the trend is decelerating.

[0037] If two points are measured, both are less
30 than the "square root of time" expectation, and the second is lower than the first, then the measured phenomenon is in quadrant III (assuming a clockwise orbit), meaning that it is in a congestion condition
35 and decelerating into deeper congestion. If two points are measured, both are less than the "square root of

09736070 121300

time" expectation, and the second is higher than the first, then the measured phenomenon is in quadrant IV (assuming a clockwise orbit), meaning that it is in a congestion condition but is accelerating out of
5 congestion toward a trend.

[0038] Finally, one could be in a situation spanning two quadrants. If two points are measured, the first exceeds the "square root of time" expectation, and the second is less than the "square root of time"
10 expectation, then the measured phenomenon has just crossed from a trend condition into a congestion condition. Similarly, if two points are measured, the first is less than the "square root of time" expectation, and the second is exceeds the "square root
15 of time" expectation, then the measured phenomenon has just crossed from a congestion condition into a trend condition.

[0039] The predictions that can be made using two points are qualitative -- e.g., "we are moving out of
20 congestion toward a trend." This is nevertheless useful. For example, if the measured phenomenon is stock price, and the determination is in quadrant IV, it may be a preferable time to buy (or sell), because while the price is about to break out into a trend, it
25 is still congested and therefore there may be willing sellers (or buyers) who do not yet realize that the price will soon turn around.

[0040] One can determine the direction of the predicted trend based on the direction of movement of
30 the actual price data at the time of transition from quadrant III to quadrant IV. That is the point where the congestion begins to decelerate. If at that point prices are moving up, the only way for the congestion to be decelerating is for the speed of price increase
35 to slow until prices start to fall, meaning the coming trend will be a downward trend. If instead at that

09736070 121300
003822 0209E760

point prices are moving down, the only way for the congestion to be decelerating is for the speed of price decrease to slow until prices start to rise, meaning the coming trend will be an upward trend.

- 5 [0041] While qualitative predictions can be made, no quantitative prediction can be made, using two points, of how long it will take, for example, to break out of congestion. Such a prediction, however, may be made using three points, equally spaced in time. Actually,
- 10 each point represents two time periods -- a shorter interval and longer interval, as discussed above, but the emphasis here is that the time between the second and third measurement points should be the same as the time between the first and second measurement points.
- 15 Because equal areas along ellipse 50 must be spanned in equal times, the area spanned between the second and third points must equal the area spanned between the first and second points. Moreover, one can then extrapolate and predict where one will be after one or
- 20 more additional equal time intervals, by determining how far along the ellipse the measurement point must move during each time interval to sweep out the same area as was swept out between the first and second points and between the second and third points.
- 25 Similarly, if one knows the time interval required to sweep a specific area, one can extrapolate and predict how long it will take to reach a specific point on the ellipse based on the total area to be swept to reach that point.
- 30 [0042] For any three orbital velocity points calculated according to the invention and evenly spaced in time, there will be only one elliptical eccentricity to which such orbital velocities can be fit so as to sweep equal areas in equal time. That ellipse can be
- 35 found using tables of elliptical motion such as are commonly used by astronomers, and which may be found,

09736070-121300

e.g., in Bauschinger, J., et al., Tafeln zur theoretischen Astronomie, 2d ed. (Leipzig, 1934).

[0043] In the discussion so far, one set of measurements has been discussed, based on a particular duration -- e.g., one-minute/four-minute measurements, or one-hour/four-hour measurements, or one-day/four-day measurements, etc. However, additional information could be gleaned by taking multiple sets of measurements. Thus, one could, at every minute, take a one-minute/four-minute measurement, and a one-hour/four-hour measurement, and a one-day/four-day measurement, etc., thereby looking at long- and short-term patterns.

[0044] Exemplary computer hardware apparatus 610 with which the present invention may be implemented is shown in FIG. 6. In FIG. 6, which shows a first preferred embodiment of apparatus according to the invention, system 610 includes a computer 611 comprising a central processing unit ("CPU") 620, a working memory 622 which may be, e.g., RAM (random-access memory) or "core" memory, mass storage memory 624 (such as one or more disk drives or CD-ROM drives), one or more cathode-ray tube ("CRT") display terminals 626, one or more keyboards 628, one or more input lines 630, and one or more output lines 640, all of which are interconnected by a conventional bidirectional system bus 650.

Input hardware 636, coupled to computer 611 by input lines 630, may be implemented in a variety of ways. Modem or modems 632, which also may be routers or other computer-to-computer communications devices, connected by a telephone line or dedicated data line (such as a T1 or T3 line) 634 can be used to allow the system to obtain real-time (or near-real-time) data, such a stock price data, from a source such as a direct feed from one of the recognized stock exchanges, or a

feed from a business data service. Modems/routers 632 also may be used to allow access through the Internet or other public access network. Alternatively or additionally, the input hardware 630 may comprise
5 CD-ROM drives or disk drives 624, particularly for input of historical data (in case a user wants to plot data over long periods such as years. In conjunction with display terminal 626, keyboard 628 may also be used as an input device, particularly to select the
10 time periods to be plotted.

Output hardware 646, coupled to computer 611 by output lines 640, may similarly be implemented by conventional devices. By way of example, output hardware 646 may include CRT display terminal 626 for
15 displaying the values of the ratios for the selected periods, or the graphical representation of those values on an ellipse if desired. Output hardware 646 might also include a printer or other printing device 642, so that hard copy output may be produced, or a
20 disk drive 624, to store system output for later use. Finally, output hardware 646 includes the necessary hardware to make the output data available to subscribers over modems or routers 632.

[0045] As stated above, implementation of the
25 present invention may be computationally intensive. For example, in a preferred embodiment, every minute the system must calculate the ratio of the range over the previous four minutes to the range over first of those four minutes, for every index and individual
30 stock or other instrument being tracked. This is in addition to similar calculations done hourly for the preceding four hour period, daily for the preceding four day period, etc. This can be achieved in a preferred embodiment by having subscribers allow their
35 computers 660 to be used, while they are connected to the system, to perform some of the calculations during

09736070-121300

25 [0048] Whether performed solely on computer 611 (or on a plurality of computers co-located with computer 611), or whether performed on a distributed basis such as on subscriber computers 660 as just described, each computer would, for a given measurement point, examine all recorded values between four (for example) time units prior to the measurement point and three time units prior to the measurement point of the price or index to be measured, and compute the difference between the maximum and minimum values during that interval. The computer would then multiply that result by two to obtain the expected range between

the longer time interval between four time units prior to the measurement point and the measurement point. The computer would then examine all recorded values of the price or index to be measured during that longer interval, and again compute the difference between the maximum and minimum values during that interval to obtain the actual range. The actual range would then be compared to the expected range as described above.

[0049] Alternatively, computer 611 or each individual subscriber computer (not shown) could be equipped with a Brownian motion standard chip as described above, which would allow a direct comparison, for any time interval, between the actual and expected range of change of the price or index to be measured. For example, the random number generator provided in the Intel® 810 chipset, from Intel Corporation, of Santa Clara, California, generates a non-deterministic "random" stream of numbers from thermal noise from a resistor. This stream of numbers, based on thermal noise, is in fact a Brownian motion standard.

[0050] The results of the data comparisons according to the invention may be displayed to users or subscribers in a number of ways. For example, as shown in FIG. 7, and particularly if the Brownian motion standard is used, the results may be displayed by graphing the expected Gaussian bell curve 70 representing a normal distribution, and then superimposing a curve 71 representing actual conditions. These curves would be for a specified time interval, such as the four-minute interval if one-minute/four-minute measurements are used, or the four-hour interval if one-hour/four-hour measurements are used. The curves would show the relative number of occurrences (each representing one transaction) of normalized price changes about a mean value, which can be expected to be zero. Thus, preferably, the peaks of

curves 70 and 71 would represent the normalized number of occurrences, or expected occurrences, of price changes of zero magnitude, and the abscissa preferably would be labelled with the number of standard
5 deviations above or below zero. The ordinate preferably would be labelled in percentages, with 100% representing the peak of Gaussian curve 70.

[0051] If the graph in FIG. 7 represents data captured during a sufficiently long interval -- e.g.,
10 four hours, the number of data points (transactions) should be sufficient to generate curve 71 with confidence. However, if there are too few data points (e.g., the interval is short, such as four minutes, and there are too few transactions), then well-known
15 conventional "bootstrapping" techniques can be used to generate a curve 71 from the available data. In such a case, there would be some uncertainty in the curve. This could be displayed to the user by, e.g., displaying a band (not shown) of a different color
20 about curve 71, with the width of the band representing the margin of error of curve 71.

[0052] Alternatively, as shown in FIG. 8, the ellipse 50 described above can be displayed, with a highlighted point 80 representing the present, and
25 points 81 representing past measurements. In addition, the eccentricity of the ellipse would reveal the relative size of the congealed group of buyers/sellers ("dissenters") as compared to the Brownian core of market participants ("majority"). For example, if the
30 ellipse is close to circular, then the number of dissenters is small and any apparent trends may be statistically insignificant, while if the ellipse is highly elongated, the number of dissenters is very large and it may be too late to join the trend (most of
35 the gains may have already occurred).

09735070.12100

5

10

25

35

moves outside band, it may signal a trend which is upward if the value is above the band or downward if the value is below the band.

[0057] Although not shown, a display integrating two
5 or more of the displays of FIGS. 7, 8 and 9 could be provided.

[0058] Thus it is seen that a model of market
activity that explains the non-normal distribution of
such activity, and a technique for market prediction
10 based on that model, has been provided. One skilled in
the art will appreciate that the present invention can
be practiced by other than the described embodiments,
which are presented for purposes of illustration and
not of limitation, and the present invention is limited
15 only by the claims which follow.

09736070 121300